

1

Supplementary Material

Assessment of observed changes and responses in natural and managed systems

Coordinating Lead Authors:

Cynthia Rosenzweig (USA), Gino Casassa (Chile)

Lead Authors:

David J. Karoly (USA/Australia), Anton Imeson (The Netherlands), Chunzhen Liu (China), Annette Menzel (Germany), Samuel Rawlins (Trinidad and Tobago), Terry L. Root (USA), Bernard Seguin (France), Piotr Tryjanowski (Poland)

Contributing Authors:

Tarekegn Abeku (Ethiopia), Isabelle Côté (Canada), Mark Dyurgerov (USA), Martin Edwards (UK), Kristie L. Ebi (USA), Nicole Estrella (Germany), Donald L. Forbes (Canada), Bernard Francou (France), Andrew Githeko (Kenya), Vivien Gornitz (USA), Wilfried Haeberli (Switzerland), John Hay (New Zealand), Anne Henshaw (USA), Terrence Hughes (Australia), Ana Iglesias (Spain), Georg Kaser (Austria), R. Sari Kovats (UK), Joseph Lam (China), Diana Liverman (UK), Dena P. MacMynowski (USA), Patricia Morellato (Brazil), Jeff T. Price (USA), Robert Muir-Wood (UK), Peter Neofotis (USA), Catherine O'Reilly (USA), Xavier Rodo (Spain), Tim Sparks (UK), Thomas Spencer (UK), David Viner (UK), Marta Vicarelli (Italy), Ellen Wiegandt (Switzerland), Qigang Wu (China), Ma Zhuguo (China)

Review Editors:

Lucka Kajfež-Bogataj (Slovenia), Jan Pretel (Czech Republic), Andrew Watkinson (UK)

This supplementary material should be cited as:

Rosenzweig, C., G. Casassa, D.J. Karoly, A. Imeson, C. Liu, A. Menzel, S. Rawlins, T.L. Root, B. Seguin and P. Tryjanowski, 2007: Supplementary material to chapter 1: Assessment of observed changes and responses in natural and managed systems. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK.

Table of Contents

1.3.2 Hydrology and water resources	SM.1-3
1.3.3 Coastal processes and zones	SM.1-4
1.3.8 Disasters and hazards	SM.1-4
1.4 Larger-scale aggregation and attribution to anthropogenic climate change	SM.1-5
Box SM.1 Linking the causes of climate change to observed effects on physical and biological systems	SM.1-7
References for Supplementary Material	SM.1-12

This Supplementary Material is in support of Chapter 1. The headings correspond to the sections of the main chapter. The Supplementary Material cannot and should not be read in isolation. It can only be read in association with the chapter.

1.3.2 Hydrology and water resources

Table SM-1.1. Examples from the literature of observed changes in (a) runoff/streamflow and (b) lake levels.

Location	Time period	Observed change	Reference
(a) Runoff/streamflow			
Arctic Ocean	1936-2002	Increasing annual discharge from six largest Eurasian rivers to Arctic Ocean by 7% due to Arctic Oscillation NAO and warming climate (this increase is not entirely consistent with apparent trends in temperature and precipitation, due to data limitations)	Peterson et al., 2002; Shiklomanov and Shiklomanov, 2003
Sweden	1807-2002	No trend in runoff due to increasing temperature offsetting the increasing precipitation	Lindstrom and Bergstrom, 2004
UK	Last 50 years	No trend in annual runoff	Hannaford and Marsh, 2005
Finland	Last 20 years	Increase in discharge	Hyvarinen, 2003
(b) Lake levels			
Bosten Lake, Xinjiang, China	1980-2000	Rise in lake level ~4 m due to increasing precipitation and snow-ice melt flow from $3.08 \times 10^8 \text{ m}^3$ to $9.6 \times 10^8 \text{ m}^3$	Yuan et al., 2003
Daihai Lake, China	1700-1996	From 1960 to 1996 lake level decreased 3.85 m due to combined effects of drought and human activities	Zhang and Ruijin, 2001
Lakes, central Italy	Last 20 years	Decline in lake level due to rainfall decrease and water withdrawal	Capelli and Mazza, 2005
Vortsjarv, Estonia	1884-2000	Strong water level fluctuation related to the North Atlantic Oscillation (NAO)	Noges et al., 2003

Table SM-1.2. Examples from the literature of observed changes in (a) floods and (b) droughts.

Location	Time period	Observed change	Reference
(a) Floods			
Global	1865-1999	Increase in frequency of floods with discharges exceeding 100-year levels from 29 large river basins more than 200,000 km ²	Milly et al., 2002
Elbe and Dresden, Germany	1997-2002	Catastrophic events much larger than 100-year flood, but no increasing trend in flood magnitude in a record from 1827	Becker and Grunewald, 2003; Kundzewicz et al., 2005
Bangladesh	1980s-1998	>50-to-100-year floods from strong monsoons	Chowdhury and Ward, 2003
Yangtze River, China	1990-1999	>50-year floods due to El Niño events	Qian and Zhu, 2001
(b) Droughts			
Much of UK	20th century	No evidence of significant increase in the occurrence of low river flows	Hannaford and Marsh, 2005
Much of Europe	1911-1995	No evidence of significant increase in droughts (defined as streamflow below a certain threshold). However, recently Europe has suffered prolonged drought associated with the severe summer heatwave	Hisdal et al., 2001; Hannaford and Marsh, 2005; van der Schrier et al., 2005
Eastern USA	1941-1999	Significant increase in annual minimum (202 out of 395 sites) and median (219 sites) daily streamflow around 1970 as a step change related to precipitation increase and NAO	Douglas, 2000; McCabe, 2002; Groisman, 2004
New England, USA	20th century	No evidence of significant increase in droughts (defined as streamflow below a certain threshold)	Hodgkins et al., 2005
Australia	2002-2003	Severe drought due to record high temperature	Karoly et al., 2003; Nicholls, 2004
Sahelian drought	1981-1993; 1994-2005	(a) 1981-1993 marked by below average Normalised Difference Vegetation Index (NDVI) and persistence of drought (b) 1994-2005 marked by a trend towards 'wetter' condition, but still far below the pre-1980s wetter condition	Hisdal et al., 2001; Hannaford and Marsh, 2005; Hodgkins et al., 2005; van der Schrier et al., 2005

Table SM-1.3. Examples from the literature of observed changes in physical and chemical water properties.


Location	Time period	Observed change	Reference
64 lakes/rivers in Europe, North America and Asia	Last 40 years	Surface water temperature warming by 0.2-1.5°C for 17 lakes. Stratified period has advanced by up to 20 days and lengthened by 2-3 weeks with increased thermal stability	Adrian and Deneke, 1996; King et al., 1998; Livingstone and Dokulil, 2001; Carvalho and Kirika, 2003; Livingstone, 2003; Straile et al., 2003; Arhonditsis et al., 2004; Dabrowski et al., 2004; Winder and Schindler, 2004
8 lakes/rivers in North America, Europe and East Africa	1991-2003 1939-2000	Decreases in nutrients in surface water and corresponding increases in deep-water concentration because of reduced upwelling due to greater thermal stability	Hambright et al., 1994; Adrian and Deneke, 1996; Straile et al., 2003
Lake Baikal, Russia	Recent decades	Decrease in silica content of 30% related to regional warming	Shimaraev et al., 2004
27 rivers, Japan	Recent decades	Increase in biological oxygen demand and suspended solids, and decrease in dissolved oxygen due to increase in air temperature	Ozaki et al., 2003

1.3.3 Coastal processes and zones

Table SM-1.4. Examples from the literature of changes in storm surges, flood heights and areas, and waves.

Type of change	Period	Location	References
More frequent and higher floods due to subsidence, hydrodynamic changes and relative sea-level rise	1830-2000	Venice, Italy	Camuffo and Stararo, 2004
Decreasing surges due to shifts in wind direction	1890s-1910s and 1950s-1997	Brittany, France	Pirazzoli et al., 2004
Increasing extreme high water levels due to climate variability and sea-level rise	1975-present	Global	Woodworth and Blackman, 2004
Decrease in mean winter significant wave height	1958-2001	Mediterranean Sea	Lionello et al., 2005

1.3.8 Disasters and hazards

**Figure SM-1.1.** An example from the literature of one study analysing rising costs of normalised weather-related catastrophes compared with global temperatures. Data smoothed over ± 4 years = 9 years until 2001 (Muir Wood et al., 2006).

1.4 Larger-scale aggregation and attribution to anthropogenic climate change

Table SM-1.5. Characteristics of the data used in the aggregation and attribution assessment of Section 1.4.

1. Database constructed of observations from studies, including information such as:
 - a. category and region (according to WGII Chapters),
 - b. longitude and latitude of study,
 - c. study dates and duration,
 - d. direction of change, consistent or not consistent with warming,
 - e. statistical significance,
 - f. type of impact and system,
 - g. whether or not land use was a driving factor.

2. Criteria for inclusion of study in synthesis assessment:
 - published peer-reviewed study,
 - statistically significant trend in change in system related to temperature or related climate variable,
 - changes observed in systems between 1970 and 2004; studies may extend after 2004,
 - studies ending in 1990 or later,
 - duration of study period 20 years or longer.




Figure SM-1.2. Duration of time-series (years) of observed changes in natural and managed systems used in statistical analysis of synthesis assessment in Section 1.4.

Table SM-1.6. Summary of observed impacts of temperature-related regional climate change in chapter synthesis assessment in Section 1.4.

Cryosphere	Changes in glaciers, lake and river ice break-up, snow cover and permafrost active layer
Hydrology	Changes in spring peak discharge and lake levels
Coastal processes	Changes in storminess and coastal vegetation, shoreline retreat, coastal erosion
Marine, freshwater and terrestrial biological systems	Changes in phenology, community composition, productivity and synchrony; shifts in latitude/altitude ranges and breeding sites; genetic adaptation

Table SM-1.7. Comparison of significant observed changes in physical and biological systems with regional temperature changes at the global scale in chapter synthesis assessment in Section 1.4.

Temperature cells	Cells with significant observed change consistent with warming*	Cells with significant observed change not consistent with warming	Cells with significant observed change consistent with warming**
Significant warming	49% (2.5%)	9% (2.5%)	56% (5%)
Warming	31% (22.5%)	4% (22.5%)	36% (45%)
Cooling	6% (22.5%)	0% (22.5%)	6% (45%)
Significant cooling	2% (2.5%)	0% (2.5%)	2% (5%)
Chi-squared value (significance level)		350 (<<1%)	104 (<<1%)

* assuming three-fold null hypothesis; ** assuming two-fold null hypothesis; see text for full explanation

Note: Fraction of $5^{\circ}\times 5^{\circ}$ cells with significant observed changes in systems (from studies considered in this chapter) and temperature changes (over 1970-2004 from HadCRUT3 – Brohan et al., 2006) in different categories (significant warming, warming, cooling, significant cooling). Expected values shown in parentheses are for the null hypotheses:

- (i) significant observed changes in systems are equally likely in each direction,
- (ii) temperature trends are due to natural climate variations and are normally distributed,
- (iii) there is no relationship between significant changes in systems and co-located warming.


The right-hand column repeats the analysis without assuming point (i) above and only considers significant observed changes in systems that are consistent with warming, in order to avoid the possible effects of publication or research biases.

The significance levels for the chi-squared values relative to the expected distribution are obtained by comparing the locations of the significant observed system changes with regional temperature trends over 35-year periods due to natural climate variability from long control simulations with 5 different coupled climate models; 192 independent 35-year periods were sampled from the control runs, allowing estimation of chi-squared values at about the 1% significance level due to natural variability.

The analysis was repeated using a second global gridded temperature dataset (GHCN-ERSST) and there were no significant differences in the results.

Footnote 1, continued from below Box SM.1 on next page. At each location, all of which are in the Northern Hemisphere, the changing trait is compared with modelled temperatures driven by: (a) Natural forcings (pink bars), (b) anthropogenic (i.e., human) forcings (orange bars), and (c) combined natural and anthropogenic forcings (yellow bars). In addition, on each panel the frequencies of the correlation coefficients between the actual temperatures recorded during each study and changes in the traits of 83 species, the only ones of the 145 with reported local-temperature trends, are shown (dark blue bars). On average the number of years species were examined is about 28 with average starting and ending years of 1960 to 1998. Note that the agreement: a) between the natural and actual plots is weaker ($K=60.16$, $P>0.05$) than b) between the anthropogenic and actual ($K=35.15$, $P>0.05$), which in turn is weaker than c) the agreement between combined and actual ($K=3.65$, $P<0.01$). Taken together, these plots show that a measurable portion of the warming regional temperatures to which species are reacting can be attributed to humans, therefore showing joint attribution (see Chapter 1).

Box SM.1. Linking the causes of climate change to observed effects on physical and biological systems. In chapter synthesis assessment in Section 1.4




The figure to the left demonstrates the linkages between observed temperatures, observed effects on natural systems, and temperatures from climate model simulations with natural, anthropogenic, and combined natural and anthropogenic forcings. Two ways in which these linkages are utilised in detection and attribution studies of observed effects are described below.

1. Using climate models


The study of causal connection by separation of natural and anthropogenic forcing factors compares observed temporal changes in animals and plants with changes over the same time periods in observed temperatures as well as modelled temperatures using (i) only natural climate forcing; (ii) only anthropogenic climate forcing; and (iii) both forcings combined.

The panel to the right shows the results from a study employing this methodology¹. The locations for the modelled temperatures were individual grid boxes corresponding to given animal and plant study sites and time periods.

The agreement (in overlap and shape) between the observed (blue bars) and modelled plots is weakest with natural forcings, stronger with anthropogenic forcings, and strongest with combined forcings. Thus, observed changes in animals and plants are likely responding to both natural and anthropogenic climate forcings, providing a direct cause-and-effect linkage [F1.7, 1.4.2.2].



2. Using spatial analysis



The study of causal connection by spatial analysis follows these stages: (i) it identifies 5° × 5° latitude/longitude cells across the globe which exhibit significant warming, warming, cooling, and significant cooling; (ii) it identifies 5° × 5° cells of significant observed changes in natural systems that are consistent with warming and that are not consistent with warming; and (iii) it statistically determines the degree of spatial agreement between the two sets of cells. In this assessment, the conclusion is that the spatial agreement is significant at the 1% level and is very unlikely to be solely due to natural variability of climate or of the natural systems.

Taken together with evidence of significant anthropogenic warming over the past 50 years averaged over each continent except Antarctica [WGI AR4² SPM], this shows a discernible human influence on changes in many natural systems [1.4.2.3].

¹ Plotted are the frequencies of the correlation coefficients (associations) between the timing of changes in traits (e.g., earlier egg-laying) of 145 species and modelled (HadCM3) spring temperatures for the grid-boxes in which each species was examined. (Continues at bottom of previous page).

² IPCC, 2007: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller, Eds., Cambridge University Press, Cambridge, 996 pp.




Figure SM-1.3. Observed changes in physical systems (cryosphere, hydrology and coastal processes) and biological systems (marine and freshwater biological systems, terrestrial biological systems) for studies ending in 1990 or later with at least 20 years of data used in chapter synthesis assessment in Section 1.4. Dots represent about 75 studies, which have >29,000 data series (of which ~27,800 are from European phenological studies of flora and fauna). Observed trends in surface air temperature and sea-surface temperature 1970-2004 (HadCRUT3 Brohan et al., 2006). White regions do not contain sufficient observational climate data to estimate a trend.




Figure SM-1.4. Changes in physical and biological systems and surface temperature used in chapter synthesis assessment in Section 1.4. Background shading, and the key to the bottom right, show changes in gridded surface temperatures over the period 1970-2004. The boxes, and the key to bottom left, show the continental-scale changes in physical (left-hand column) and biological (right-hand column) systems calculated from individual series with at least 20 years data in the 1970-2004 period; the top row shows the number of observed series matching the length criterion that show a significant trend and the bottom row shows the percentage of these in which the trend is consistent with warming. At the global scale TER = Terrestrial; MFW = Marine and Freshwater, and GLO = Global.

Table SM-1.8. References of studies that fit the criteria selected for global synthesis assessment in Section 1.4.**Database Reference List**

Year	Short reference	Full reference
1 2001	Abu-Asab et al. (2001)	Abu-Asab, et al., 2001: Earlier plant flowering in spring as a response to global warming in the Washington, DC, area. <i>Biodiversity and Conservation</i> , 10 , 597-612.
2 2002	Ahas et al. (2002)	Ahas, R., et al., 2002: Changes in European spring phenology. <i>International Journal of Climatology</i> , 22 , 1727-1738.
3 2006	Allan and Komar (2006)	Allan, J. C. and P. D. Komar, 2006: Climate controls on U.S. west coast erosion processes. <i>Journal of Coastal Research</i> , 3 , 511-529.
4 2002	Arendt et al. (2002)	Arendt, A.A., et al., 2002: Rapid wastage of Alaska glaciers and their contribution to rising sea level. <i>Science</i> , 297 , 382-386.
5 2004	Atkinson et al. (2004)	Atkinson, A., et al., 2004: Long term decline in krill stock and increase in salps within the Southern Ocean. <i>Nature</i> , 432 , 100-103.
6 2001	Barbraud and Weimerskirch (2001)	Barbraud C. and H. Weimerskirch, 2001: Emperor penguins and climate change. <i>Nature</i> , 411 , 183-186.
7 1995	Barry et al. (1995)	Barry, J.P., et al., 1995: Climate-related, long-term faunal changes in a California rocky intertidal community. <i>Science</i> , 267 , 672-965.
8 2000	Beaubien and Freeland (2000)	Beaubien, E.G. and H.J. Freeland, 2000: Spring phenology trends in Alberta, Canada: links to ocean temperature. <i>International Journal of Biometeorology</i> , 44 , 53-59.
9 2002	Beaugrand et al. (2002)	Beaugrand, G., et al., 2002: Reorganization of North Atlantic marine copepod biodiversity and climate. <i>Science</i> , 296 , 1692-1694.
10 2004	Both et al. (2004)	Both, C., et al., 2004: Large-scale geographical variation confirms that climate change causes birds to lay earlier. <i>Proceedings of the Royal Society of London Series B – Biological Sciences</i> , 271 , 1657.
11 2001	Bradshaw, and Holzapfel (2001)	Bradshaw, W.E. and C.M. Holzapfel, 2001: Genetic shift in photoperiodic response correlated with global warming. <i>Proceedings of the National Academy of Sciences of the USA</i> , 98 , 14509.
12 2004	Brooks and Birks (2004)	Brooks, S.J. and H.J.B. Birks, 2004: The dynamics of Chironomidae (Insecta: Diptera) assemblages in response to environmental change during the past 700 years on Svalbard. <i>Journal of Paleolimnology</i> , 31 , 483-498.
13 1999 (cited in TAR)	Brown et al. (1999)	Brown, J.L., et al., 1999: Long-term trend toward earlier breeding in an American bird: a response to global warming? <i>Proceedings of the National Academy of Sciences of the USA</i> , 96 , 5565-5569.
14 2002	Bunce et al. (2002)	Bunce, A., et al., 2002: Long-term trends in the Australasian gannet (<i>Morus serrator</i>) population in Australia: the effect of climate change and commercial fisheries. <i>Marine Biology</i> , 141 , 263-269.
15 2003	Butler (2003)	Butler, C.J., 2003: The disproportionate effect of global warming on the arrival dates of short-distance migratory birds in North America. <i>Ibis</i> , 45 , 484.
16 2005	Chambers (2005)	Chambers, L.E., 2005: Migration dates at Eyre Bird Observatory: links with climate change? <i>Climate Research</i> , 29 , 157-165.
17 2004	Chuine et al. (2004)	Chuine, I., et al., 2004: Grape ripening as a past climate indicator. <i>Nature</i> , 432 , 289-290.
18 2005	Cook et al. (2005)	Cook, A.J., et al., 2005: Retreating glacier fronts on the Antarctic Peninsula over the past half-century. <i>Science</i> , 308 , 541-544.
19 2004	Corn (2003)	Corn, P.S., 2003: Amphibian breeding and climate change: importance of snow in the mountains. <i>Conservation Biology</i> , 17 , 622-625.
20 2001	Dafila and Clot (2001)	Dafila, C. and B. Clot, 2001: Phytophenological trends in Switzerland. <i>International Journal of Biometeorology</i> , 45 , 203-207.
21 2001	D'Arrigo et al. (2001)	D'Arrigo, R., et al., 2001: 1738 years of Mongolian temperature variability inferred from tree-ring chronology of Siberian pine. <i>Geophysical Research Letters</i> , 28 , 543-546.
22 2003	Daufresne et al. (2003)	Daufresne, M., et al., 2004: Long-term changes within the invertebrate and fish communities of the Upper Rhone River: effects of climatic factors. <i>Global Change Biology</i> , 10 , 124-140.
23 2005	Dyurgerov and Meier (2005)	Dyurgerov, M.B. and M.F. Meier, 2005: Glaciers and the changing earth system: a 2004 snapshot. Occasional Paper No. 58. INSTAAR, University of Colorado at Boulder, Colorado.
24 2004	Edwards and Richardson (2004)	Edwards, M. and A.J. Richardson, 2004: Impact of climate change on marine pelagic phenology and trophic mismatch. <i>Nature</i> , 430 , 881- 884.
25 2006	Field et al. (2006)	Field, D.B., et al., 2006: Planktonic foraminifera of the California current reflect 20th-century warming. <i>Science</i> , 311 , 63-66.
26 2002	Fitter and Fitter (2002)	Fitter, A. H. and R.S.R. Fitter, 2002: Rapid changes in flowering time in British plants. <i>Science</i> , 296 , 1689-1691.
27 2004	Forbes et al. (2004)	Forbes, D.L., et al., 2004: Storms and shoreline retreat in the southern Gulf of St. Lawrence. <i>Marine Geology</i> , 210 , 169-204.
28 2003	Forister and Shapiro (2003)	Forister, M.L. and A.M. Shapiro, 2003: Climatic trends and advancing spring flight of butterflies in lowland California. <i>Global Change Biology</i> , 9 , 1130-1135.
29 2004	Frauenfeld et al. (2004)	Frauenfeld, O.W., et al., 2004: Interdecadal changes in seasonal freeze and thaw depths in Russia. <i>Journal of Geophysical Research</i> , 109 , D5101.
30 2004	Georges (2004)	Georges, C., 2004: 20th-century glacier fluctuations in the tropical Cordillera Blanca, Peru. <i>Arctic, Antarctic and Alpine Research</i> , 36 , 100-107.
31 2001	Gibbs and Breish (2001)	Gibbs, J.P. and A.R. Breish, 2001: Climate warming and calling phenology of frogs near Ithaca New York, 1900-1999. <i>Conservation Biology</i> , 15 , 1175-1178.
32 2005	Hampton (2005)	Hampton, S.E., 2005: Increased niche differentiation between two <i>Conochilus</i> species over 33 years of climate change and food web alteration. <i>Limnology and Oceanography</i> , 50 , 421-426.

- 33 2003 Hennessy et al. (2003) Hennessy, K., et al., 2003: *The Impact of Climate Change on Snow Conditions in Mainland Australia*. CSIRO, Aspendale.
- 34 2003 Hodgkins et al. (2003) Hodgkins, G.A., et al., 2003: Changes in the timing of high river flows in New England over the 20th century. *Journal of Hydrology*, **278**, 244-252.
- 35 2000 Inouye et al. (2000) Inouye, D.W., et al., 2000: Climate change is affecting altitudinal migrants and hibernating species. *Proceedings of the National Academy of Sciences of the USA*, **97**, 1630.
- 36 2002 Jacobs et al. (2002) Jacobs, S.S., et al., 2002: Freshening of the Ross Sea during the late 20th century. *Science*, **297**, 386-389.
- 37 2001 Jones et al. (2001) Jones, R.N., et al., 2001: Modelling historical lake levels and recent climate change at three closed lakes, Victoria, Australia (c. 1840-1990). *Journal of Hydrology*, **246**, 159-180.
- 38 2004 Juanes et al. (2004) Juanes, F., et al., 2004: Long-term changes in migration timing of adult Atlantic salmon (*Salmo salar*) at the southern edge of the species distribution. *Canadian Journal of Fisheries and Aquatic Sciences*, **61**, 2392-2400.
- 39 2005 Karst-Riddoch et al. (2005) Karst-Riddoch, T.L., et al., 2005: Diatom responses to 20th century climate-related environmental changes in high-elevation mountain lakes of the northern Canadian Cordillera. *Journal of Paleolimnology*, **33**, 265-282.
- 40 2002 Kozlov and Berlina (2002) Kozlov, M.V. and N.G. Berlina, 2002: Decline in length of the summer season in the Kola peninsula, Russia. *Climatic Change*, **54**, 387-398.
- 41 2002 Kullman (2002) Kullman, L., 2002: Rapid recent range-margin rise of tree and shrub species in the Swedish Scandes. *Journal of Ecology*, **90**, 68.
- 42 2004 Ledneva et al. (2004) Ledneva, A., et al., 2004: Climate change as reflected in a naturalist's diary, Middleborough, Massachusetts. *Wilson Bulletin*, **116**, 224-231.
- 43 2004 Lips et al. (2004) Lips, K.R., et al., 2004: Amphibian population declines in montane southern Mexico: resurveys of historical localities. *Biological Conservation*, **119**, 555-564.
- 44 2000 (Cited Magnuson et al. (2000) Magnuson, J. J., et al., 2000: Historical trends in lake and river ice cover in the Northern Hemisphere. *Science*, **289**, 1743-1746.
- in TAR)
- 45 2003 Matsumoto et al. (2003) Matsumoto, K., et al., 2003: Climate change and extension of the Ginko biloba L. growing season in Japan. *Global Change Biology*, **9**, 1634-1642.
- 46 2006 Menzel et al. (2006) Menzel, A., et al., 2006: European phenological response to climate change matches the warming pattern. *Global Change Biology*, **12**, 1969-1976.
- 47 2003 Michelutti et al. (2003) Michelutti, N., et al., 2003: Diatom response to recent climatic change in a high arctic lake (Char Lake, Cornwallis Island, Nunavut). *Global and Planetary Change*, **38**, 257-271.
- 48 2005 Mills (2005) Mills, A.M., 2005: Changes in the timing of spring and autumn migration in North American migrant passerines during a period of global warming. *Ibis*, **147**, 259-269.
- 49 2005 Mote et al. (2005) Mote, P.W., et al., 2005: Declining mountain snowpack in western North America. *Bulletin of the American Meteorological Society*, January, 39-49.
- 50 2003 O'Reilly et al. (2003) O'Reilly, C.M., et al., 2003: Climate change decreases aquatic ecosystem productivity of Lake Tanganika, Africa. *Nature*, **424**, 766-768.
- 51 2005 Oerlemans (2005) Oerlemans, J., 2005: Extracting a climate signal from 169 glacial records. *Science*, **208**, 675-677.
- 52 2003 Orviku et al. (2003) Orviku, K., et al., 2003: Increasing activity of coastal processes associated with climate change in Estonia. *Journal of Coastal Research*, **19**, 364-375.
- 53 2003 Peñuelas and Boada (2003) Peñuelas, J. and M. Boada, 2003: A global change-induced biome shift in the Montseny Mountains (NE Spain). *Global Change Biology*, **9**, 131-140.
- 54 2002 Peñuelas et al. (2002) Peñuelas, J., et al., 2002: Changed plant and animal life cycles from 1952 to 2000 in the Mediterranean region. *Global Change Biology*, **8**, 531-544.
- 55 2003 Perren et al. (2003) Perren, B.B., et al., 2003: Rapid lacustrine response to recent High Arctic warming: a diatom record from Sawtooth Lake, Ellesmere Island, Nunavut. *Arctic, Antarctic and Alpine Research*, **35**, 271-278.
- 56 2004 Primack et al. (2004) Primack, D., et al., 2004: Herbarium specimens demonstrate earlier flowering times in response to warming in Boston. *American Journal of Botany*, **91**, 1260-1264.
- 57 2004 Punsalma et al. (2004) Punsalma, B., et al., 2004: Trends in river and lake ice in Mongolia. Working Paper No. 4. AIACC, http://www.aiaccproject.org/working_papers/Working%20Papers/AIACC_WP_No004.pdf.
- 58 2004 Richardson and Schoeman (2004) Richardson, A.J. and D.S. Schoeman, 2004: Climate impacts on plankton ecosystems in the northeast Atlantic. *Science*, **305**, 1609-1612.
- 59 2003 Rignot et al. (2003) Rignot, E., et al., 2003: Contribution of the Patagonia icefields of South America to sea level rise. *Science*, **302**, 434-437.
- 60 2000 Roetzer et al. (2000) Roetzer, T., et al., 2000: Phenology in central Europe: differences and trends of spring phenophases in urban and rural areas. *International Journal of Biometeorology*, **44**, 60-66.
- 61 2003 Ron et al. (2003) Ron, S.R., et al., 2003: Population decline of the Jambato toad *Atelopus ignescens* (Anura: Bufonidae) in the Andes of Ecuador. *Journal of Herpetology*, **37**, 116-126.
- 62 2002 Sorvari et al. (2002) Sorvari, S., et al., 2002: Lake diatom response to recent Arctic warming in Finnish Lapland. *Global Change Biology*, **8**, 171-181.
- 63 2005 Stervander et al. (2005) Stervander, M., et al., 2005: Timing of spring migration in birds: long-term trends, North Atlantic Oscillation and the significance of different migration routes. *Journal of Avian Biology*, **36**, 210-221.
- 64 2004 Sumner et al. (2004) Sumner, P.D., et al., 2004: Climate change melts Marion Island's snow and ice. *South African Journal of Science*, **100**, 395-398.
- 65 2002 Van Duivenbooden et al. (2002) Van Duivenbooden, N., et al., 2002: Impact of climate change on agricultural production in the Sahel. Part 2. Case study for groundnut and cowpea in Niger. *Climatic Change*, **54**, 349-368.

- 66 2002 van Vliet et al. (2002) van Vliet, A.J.H., et al., 2002: The influence of temperature and climate change on the timing of pollen release in the Netherlands. *International Journal of Climatology*, **22**, 1757-1767.
- 67 2003 Visser et al. (2003) Visser, M.E., et al., 2003: Variable responses to large-scale climate change in European Parus populations. *Proceedings of the Royal Society of London Series B – Biological Sciences*, **270**, 367-372.
- 68 2004 Williams and Abberton (2004) Williams, T.A. and M.T. Abberton, 2004: Earlier flowering between 1962 and 2002 in agricultural varieties of white clover. *Oecologia*, **138**, 122-126.
- 69 2001 Wolfe and Perren (2001) Wolfe, A.P. and B.B. Perren, 2001: Chrysophyte microfossils record marked responses to recent environmental changes in high- and mid-arctic lakes. *Canadian Journal of Botany – Revue Canadienne de Botanique*, **79**, 747-752.
- 70 2005 Wolfe et al. (2005) Wolfe, D.W., et al., 2005: Climate change and shifts in spring phenology of three horticultural woody perennials in northeastern USA. *International Journal of Biometeorology*, **49**, 303-309.
- 71 2001 Yom-Tov (2001) Yom-Tov, Y., 2001: Global warming and body mass decline in Israeli passerine birds. *Proceedings of the Royal Society of London Series B – Biological Sciences*, **268**, 947.
- 72 2002 Yoo and D'Odorico (2002) Yoo, J.C. and P. D'Odorico, 2002: Trends and fluctuations in the dates of ice-break-up of lakes and rivers in Northern Europe: the effect of the Northern Atlantic Oscillation. *Journal of Hydrology*, **268**, 100-112.
- 73 2003 Yoshikawa (2003) Yoshikawa, K. and L.D. Hinzman, 2003: Shrinking thermokarst ponds and groundwater dynamics in discontinuous and Hinzman permafrost. *Permafrost and Periglacial Processes*, **14**, 151-160.

References for Supplementary Material

- Adrian, R. and R. Deneke, 1996: Possible impact of mild winters on zooplankton succession in eutrophic lakes of the Atlantic European area. *Freshwater Biology*, **36**, 757-770.
- Anyamba, A. and Tucker, C.J., 2005: Analysis of Sahelian vegetation dynamics using NOAA-AVHRR NDVI data from 1981-2003. *J. Arid Environments*, **63**, 596-614.
- Arhonditsis, G.B., M.T. Brett, C.L. DeGasperi and D.E. Schindler, 2004: Effects of climatic variability on the thermal properties of Lake Washington. *Limnology and Oceanography*, **49**, 256-270.
- Becker, A. and U. Grunewald, 2003: Disaster management: flood risk in central Europe. *Science*, **300**, 1099-1099.
- Brohan, P., J.J. Kennedy, I. Harris, S.F.B. Tett and P.D. Jones, 2006: Uncertainty estimates in regional and global observed temperature changes: a new data set from 1850. *Journal of Geophysical Research*, **111**, D12106, doi:10.1029/2005JD006548.
- Camuffo, D. and G. Stararo, 2004: Use of proxy-documentary and instrumental data to assess the risk factors leading to sea flooding in Venice. *Global and Planetary Change*, **40**, 93-103.
- Capelli, G. and R. Mazza, 2005: Water criticality in the Colli Albani (Rome, Italy). *Giornale di Geologia Applicata*, **2005**, 113-121.
- Carvalho, L. and A. Kirika, 2003: Changes in shallow lake functioning: response to climate change and nutrient reduction. *Hydrobiologia*, **506**, 789-796.
- Chowdhury, M.R. and M.N. Ward, 2003: Seasonal rainfall and streamflow in the Ganges–Brahmaputra basins of Bangladesh: variability and predictability. *ASCE Conference Proceedings*, **118**, 29.
- Dabrowski, M., W. Marszelewski and R. Skowron, 2004: The trends and dependencies between air and water temperatures in lakes in northern Poland from 1961-2000. *Hydrology and Earth System Sciences*, **8**, 79-87.
- Douglas, E.M., R.M. Vogel and C.N. Kroll, 2000: Trends in floods and low flows in the United States: Impacts of spatial correlation. *Journal of Hydrology*, **240**, 90-105.
- Groisman, P.Y., R.W. Knight, T.R. Karl and Co-authors, 2004: Contemporary changes of the hydrological cycle over the contiguous United States: trends derived from in situ observations. *Journal of Hydrometeorology*, **5**, 64-85.
- Hambright, K.D., M. Gophen and S. Serruya, 1994: Influence of long-term climatic changes on the stratification of a subtropical, warm monomictic lake. *Limnology and Oceanography*, **39**, 1233-1242.
- Hannaford, J. and T.J. Marsh, 2005: An assessment of trend in UK runoff and low flows using a network of undisturbed catchments. *International Journal of Climatology*, **26**, 1237-1253.
- Herrmann, S. M., A. Anyamba, and C.J. Tucker, 2005: Recent trends in vegetation dynamics in the African Sahel and their relationship to climate. *Global Environmental Change*, **15**, 394-404.
- Hisdal, H., K. Stahl, L.M. Tallaksen and S. Demuth, 2001: Have streamflow droughts in Europe become more severe or frequent? *International Journal of Climatology*, **21**, 317-333.
- Hodgkins, G.A., R.W. Dudley and T.G. Huntington, 2005: Changes in the number and timing of days of ice-affected flow on northern New England rivers, 1930-2000. *Climatic Change*, **71**, 319-340.
- Hodgkins, G.A., R.W. Dudley, and T.G. Huntington, 2005: Summer low flows in New England during the 20th century. *Journal of the American Water Resources Association*, **41**, 403-412.
- Hyvarinen, V., 2003: Trend and characteristics of hydrological time series in Finland. *Nordic Hydrology*, **34**, 71-91.
- Karoly, D.J. and Co-authors, 2003: Global warming contributes to Australia's worst drought. *Australasian Science*, April, 14-17.
- King, J.R., B.J. Shuter and A.P. Zimmerman, 1998: The response of the thermal stratification of South Bay (Lake Huron) to climatic variability (vol 54, pg 1873, 1997). *Canadian Journal of Fisheries and Aquatic Sciences*, **55**, 1999-1999.
- Kundzewicz, Z.W., U. Ulbrich, T. Brucher, D. Graczyk, A. Kruger, G.C. Leckebusch, L. Menzel, I. Pinskiwar, M. Radziejewski and M. Szwed, 2005: Summer floods in central Europe: climate change track? *Natural Hazards*, **36**, 165-189.
- Lindstrom, G. and S. Bergstrom, 2004: Runoff trends in Sweden 1807-2002. *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques*, **49**, 69-83.
- Lionello, P., J. Bhend, A. Buzzi, P.M. Della-Marta, S. Krichak, A. Jansa, P. Maheras, A. Sanna, I.F. Trigo and R. Trigo, 2005: Cyclones in the Mediterranean region: climatology and effects on the environment. *Mediterranean Climate Variability*, P. Lionello, P. Malanotte-Rizzoli and R. Boscolo, Eds., Elsevier, Amsterdam, 324-372.
- Livingstone, D.M., 2003: Impact of secular climate change on the thermal structure of a large temperate central European lake. *Climatic Change*, **57**, 205-225.
- Livingstone, D.M. and M.T. Dokulil, 2001: Eighty years of spatially coherent Austrian lake surface temperatures and their relationship to regional air temperature and the North Atlantic Oscillation. *Limnology and Oceanography*, **46**, 1220-1227.
- McCabe, G.J. and D.M. Wolock, 2002: A step increase in streamflow in the conterminous United States. *Geophysical Research Letters*, **29**, 2185.
- Milly, P.C.D., R.T. Wetherald, K.A. Dunne and T.L. Delworth, 2002: Increasing risk of great floods in a changing climate. *Nature*, **415**, 514-517.
- Muir Wood R., S. Miller and A. Boissonnade, 2006: The search for trends

- in global catastrophe losses. Workshop on Climate Change and Disaster Losses: Understanding and Attributing Trends and Projections. May 25-26, 2006. Hohenkammer, Munich. Final Workshop Report, 188-194.
- Nicholls, N., 2004: The changing nature of Australian droughts. *Climate Change*, **63**, 323-336.
- Nicholson, S., 2005: On the question of the "recovery" of the rains in the West African Sahel. *Journal of Arid Environments*, **63**, 615-641.
- Noges, T., P. Noges and R. Laugaste, 2003: Water level as the mediator between climate change and phytoplankton composition in a large shallow temperate lake. *Hydrobiologia*, **506**, 257-263.
- Olsson, L., L. Eklundh and J. Ardoe, 2005: A recent greening of the Sahel-trends, patterns and potential causes. *Journal of Arid Environments*, **63**, 556-566.
- Ozaki, N., T. Fukushima, H. Harasawa, T. Kojiri, K. Kawashima and M. Ono, 2003: Statistical analyses on the effects of air temperature fluctuations on river water qualities. *Hydrological Processes*, **17**, 2837-2853.
- Peterson, B.J., R.M. Holmes, J.W. McClelland, C.J. Vorosmarty, R.B. Lammers, A.I. Shiklomanov, I.A. Shiklomanov and S. Rahmstorf, 2002: Increasing river discharge to the Arctic Ocean. *Science*, **298**, 2171-2173.
- Pirazzoli, P.A., H. Regnaud and L. Lemasson, 2004: Changes in storminess and surges in western France during the last century. *Marine Geology*, **210**, 307-323.
- Qian, W.H. and Y.F. Zhu, 2001: Climate change in China from 1880 to 1998 and its impact on the environmental condition. *Climatic Change*, **50**, 419-444.
- Shiklomanov, I.A. and A.I. Shiklomanov, 2003: Climate change and dynamics of freshwater inflow to the Arctic Ocean (in Russian). *Vodnye Resursy*, **30**, 645-654.
- Shimaraev, M.N., V.M. Domysheva and I.B. Mizandrontsev, 2004: Climate and processes in Lake Baikal ecosystem at present (in Russian). *The Vith All-Russia Hydrological Congress, Abstracts of Papers*, **4**, 287-288.
- Straile, D., K. Johns and H. Rossknecht, 2003: Complex effects of winter warming on the physicochemical characteristics of a deep lake. *Limnology and Oceanography*, **48**, 1432-1438.
- van der Schrier, G., K.R. Briffa, P.D. Jones and T.J. Osborn, 2006: Summer moisture variability across Europe. *Journal of Climate*, **19**, 2818-2834.
- Winder, M. and D.E. Schindler, 2004: Climate change uncouples trophic interactions in an aquatic system. *Ecology*, **85**, 3178-3178.
- Woodworth, P.L. and D.L. Blackman, 2004: Evidence for systematic changes in extreme high waters since the mid-1970s. *Journal of Climate*, **17**, 1190-1197.
- Yuan, L., T. Yuan, J. Fengqing, W. Yajun and H. Ruiji, 2003: Study on hydrological features of the Kaidu River and the Bosten Lake in the second half of 20th century (in Chinese). *Journal of Glaciology and Geocryology*, **25**, 215-217.
- Zhang, Z. and W. Ruijin, 2001: Climatic variation and human activities recorded by lake sediment in the catchment of Daihai Lake during the past 300 years (in Chinese). *Journal of Capital Normal University (Natural Science Edition)*, **22**, 70-76.